

Treprostinil Injection is available through the Specialty Pharmacy (SP) provider listed on page 7.

Complete all sections on this enrollment form. Let your patient know that the Specialty Pharmacy will be calling to process their prescription and that it is important to answer or return any messages.

Sign the Statement of Medical Necessity on page 3 for the Prescription.

Sign at the bottom of pages 4 and 5.

Fax the enrollment form and signed supporting documents (use Fax Cover Sheet provided on page 7) to the SP.

Information regarding the Centers for Medicare and Medicaid Services (CMS) established and expected coverage criteria for prostacyclin is included for your convenience.

MEDICARE COVERAGE CRITERIA FOR PROSTACYCLIN

The current Local Coverage Determination for Prostacyclin is as follows:

The pulmonary hypertension is not secondary to pulmonary venous hypertension (e.g., left sided atrial or ventricular disease, left sided valvular heart disease, etc.) or disorders of the respiratory system (e.g., chronic obstructive pulmonary disease, interstitial lung disease, obstructive sleep apnea or other sleep disordered breathing, alveolar hypoventilation disorders, etc.); and

The patient has idiopathic/heritable pulmonary hypertension or pulmonary hypertension which is associated with one of the following conditions: connective tissue disease, thromboembolic disease of the pulmonary arteries, human immunodeficiency virus (HIV) infection, cirrhosis, diet drugs, congenital left to right shunts, etc.

If the above conditions are present, the following criteria must be met:

- The pulmonary hypertension has progressed despite maximal medical and/or surgical treatment of the identified condition; and
- The mean pulmonary artery pressure is greater than 25 mm Hg at rest or greater than 30 mm Hg with exertion; and
- The patient has significant symptoms from the pulmonary hypertension (i.e., severe dyspnea on exertion, and either fatigability, angina, or syncope); and
- Treatment with oral calcium channel blocking agents has been tried and failed or has been considered and ruled out.

Medicare coverage criteria provided for informational purposes only. Please check with the payer to verify billing requirements. Liquidia and Sandoz do not make any representation or guarantees concerning reimbursement or coverage for any service or item.

PATIENT INFORMATION

Patient Name (first, MI, last)

Address

City

State

Zip

SHIPPING ADDRESS (if different from above):

Address

City

State

Zip

CAREGIVER

Caregiver Name

Caregiver Email

Date of Birth (mm/dd/yyyy)

Gender: _____

Email

Phone

- Home
 Cell
 Other

Alternate Phone

- Home
 Cell
 Other

Preferred contact: Phone Email

Best time to call: Morning Afternoon Night

OK to leave message with Caregiver? Yes No

INSURANCE INFORMATION

Pharmacy Benefits Manager

PRIMARY Medical Insurance Carrier

Policyholder Name

Policy ID Number

Group No (if applicable)

Medical Insurance Phone

Relationship to Policyholder

Please include copies of the front and back of all patient's medical and prescription insurance cards.

SECONDARY Medical Insurance Carrier

Policyholder Name

Policy ID Number

Group No (if applicable)

Medical Insurance Phone

Relationship to Policyholder

<input type="text"/>	<input type="text"/>
Patient Name (first, MI, last)	Date of Birth

PRESCRIBER INFORMATION

<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Prescriber Name (first, MI, last)	NPI #	State License #	Tax ID #
<input type="text"/>	<input type="text"/>		
Office/Clinic/Institution Name	Office Contact Name		
<input type="text"/>	<input type="text"/>		
Address	Office Contact Email		
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
City	State	Zip	Phone
			Fax
Preferred method of communication: <input type="radio"/> Phone <input type="radio"/> Email <input type="radio"/> Fax			

PRESCRIPTION INFORMATION

Sandoz® Treprostinil Injection vial concentration

NDC(s) prescribed:

- 1 mg/mL (20-mL vial) (00781-3420-80)
- 2.5 mg/mL (20-mL vial) (00781-3425-80)
- 5 mg/mL (20-mL vial) (00781-3427-80)
- 10 mg/mL (20-mL vial) (00781-3430-80)

Diluent: (0.9% Sodium Chloride will be used if no box is checked)

- 0.9% Sodium Chloride for Injection
- Sandoz® Sterile Diluent for Treprostinil Injection
- Sterile Water for Injection
- Epoprostenol Sterile Diluent for Injection

Infusion route and pumps:

- Subcutaneous continuous infusion with appropriate ambulatory infusion pump.
- Intravenous continuous infusion with appropriate ambulatory infusion pump.

Dosing and titration instructions

Patient dosing weight:

 kg lb

Initiation dosage:

 ng/kg/min

Titrate by _____ ng/kg/min every _____ days until goal of _____ ng/kg/min is achieved.

Indicate any alternative or additional titration instructions here:

- Dispense 1 month of drug, needles, syringes, ancillary supplies, and medical equipment necessary to administer medication. _____ refills

STATEMENT OF MEDICAL NECESSITY

Prescriber Signature is Required to Validate Prescriptions.

I certify that the pulmonary arterial hypertension therapy ordered above is medically necessary and that I am personally supervising the care of this patient.

Prescriber Full Name (print)

Dispense As Written (DAW) / Brand Medically Necessary / No Substitution / May Not Substitute / Do Not Substitute

Substitution Permitted / May Substitute / Product Selection Permitted

SIGN HERE

Prescriber Signature*

Prescriber Signature*

Date

CA, MA, NC & PR: Interchange is mandated unless Prescriber writes the words "No Substitution": _____

The prescriber is to comply with his/her state specific prescription requirements such as e-prescribing, state specific prescription form, fax language, etc. Non-compliance with state specific requirements could result in outreach to the prescriber.

*Prescriber attests that this is his/her legal signature.

No Stamps. Prescriptions Must Be Faxed.

NOTE: The responsibility to determine coverage and reimbursement parameters, and appropriate coding for a particular patient and/or procedure, is the responsibility of the provider. The information provided here is not a guarantee of coverage or reimbursement.

<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Patient Name (first, MI, last)	Date of Birth	Prescriber Name (first, MI, last)	NPI #

NURSING ORDERS

NURSE VISITS (select **one** option)

- SP home healthcare RN visit(s) to provide assessment and education on self-administration of Treprostinil to include dose, titration, and side effect management **OR**
- Prescriber-directed SP home healthcare RN visit(s) as detailed below:

Location: Home Outpatient clinic Hospital Virtual

SITE CARE

- Dressing change every _____ days
- Per standard of care

CALCIUM CHANNEL BLOCKER STATEMENT

Indicate whether the patient named above was trialed on a calcium channel blocker prior to the initiation of therapy and provide the results.

A calcium channel blocker was not trialed because:

- Patient has depressed cardiac input
- Patient has systematic hypotension
- Patient has known hypersensitivity
- Patient is hemodynamically unstable or has a history of postural hypotension
- Patient did not meet ACCP Guidelines for Vasodilator Response
- Patient has documented brachycardia or second or third-degree heartblock
- Other: _____

The following calcium channel blocker was trialed:

The patient had the following response(s):

- Patient hypersensitive or allergic
- Adverse event
- Patient became hemodynamically unstable
- Pulmonary arterial pressure continued to rise
- Disease continued to progress, or patient remained symptomatic
- Other: _____

PRESCRIBER SIGNATURE

**SIGN
HERE**

Prescriber Full Name (print)

Prescriber Signature

Date

NOTE: The responsibility to determine coverage and reimbursement parameters, and appropriate coding for a particular patient and/or procedure, is the responsibility of the provider. The information provided here is not a guarantee of coverage or reimbursement.

Patient Name (first, MI, last)	Date of Birth	Prescriber Name (first, MI, last)	NPI #

PATIENT EVALUATION

Patient Status:
 Outpatient
 Inpatient

Treprostinil Injection Status:
 Naïve / New
 Restart
 Transition

Allergies:
 No known drug allergies (NKDA)
 Yes (specify): _____

Diabetic?
 Yes
 No

WHO Group: _____

NYHA Functional Class: I II III IV

Current Medications (list all):

Date Taken:

Weight: kg lb

Height: cm in

MEDICAL INFORMATION

REQUIRED: Please select one of the following ICD-10 codes, or Other ICD-10 code, as applicable. The following ICD-10 codes do not suggest approval, coverage, or reimbursement for specific uses or indications.

DIAGNOSIS

ICD-10 I27.0 Primary pulmonary hypertension
 Idiopathic PAH
 Heritable PAH

ICD-10 I27.2 Other secondary pulmonary hypertension
 Connective tissue disease
 Drugs/Toxins induced
 HIV
 Congenital heart disease
 Portal hypertension

Other ICD-10:
 Code Description

TREATMENT HISTORY

Please indicate treatment history

Adempas® (riociguat) Tablets	<input type="radio"/> Current	<input type="radio"/> Discontinued
Epoprostenol Sodium for Injection	<input type="radio"/> Current	<input type="radio"/> Discontinued
Flolan® (epoprostenol sodium) for Injection	<input type="radio"/> Current	<input type="radio"/> Discontinued
Letairis® (ambrisentan) Tablets	<input type="radio"/> Current	<input type="radio"/> Discontinued
Opsumit® (macitentan) Tablets	<input type="radio"/> Current	<input type="radio"/> Discontinued
Orenitram® (treprostinil) Extended-Release Tablets	<input type="radio"/> Current	<input type="radio"/> Discontinued
PDE-5i (specify drugs):	<input type="radio"/> Current	<input type="radio"/> Discontinued
Remodulin® (treprostinil) Injection	<input type="radio"/> Current	<input type="radio"/> Discontinued
Tracleer® (bosentan) Tablets	<input type="radio"/> Current	<input type="radio"/> Discontinued
Tyvaso® (treprostinil) Inhalation Solution	<input type="radio"/> Current	<input type="radio"/> Discontinued
Tyvaso® DPI (treprostinil) Inhalation Powder	<input type="radio"/> Current	<input type="radio"/> Discontinued
Uptravi® (selexipag) Tablets	<input type="radio"/> Current	<input type="radio"/> Discontinued
Velettri® (epoprostenol) for Injection	<input type="radio"/> Current	<input type="radio"/> Discontinued
Ventavis® (iloprost) Inhalation Solution	<input type="radio"/> Current	<input type="radio"/> Discontinued
Other: _____	<input type="radio"/> Current	<input type="radio"/> Discontinued

TRANSITION STATEMENT (if applicable)

It is necessary for this patient to transition

from:

to:

Please provide justification for this transition.

PRESCRIBER SIGNATURE

SIGN HERE

Prescriber Full Name (print)	Prescriber Signature	Date

NOTE: The responsibility to determine coverage and reimbursement parameters, and appropriate coding for a particular patient and/or procedure, is the responsibility of the provider. The information provided here is not a guarantee of coverage or reimbursement.

INDICATION

Treprostinil injection is a prostacyclin mimetic indicated for

- Treatment of pulmonary arterial hypertension (PAH), World Health Organization (WHO) Group 1, to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with New York Heart Association (NYHA) Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%).
- Patients who require transition from epoprostenol to reduce the rate of clinical deterioration. The risks and benefits of each drug should be carefully considered prior to transition.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

None.

WARNINGS AND PRECAUTIONS

- Chronic intravenous (IV) infusions delivered using an external infusion pump with an indwelling central venous catheter are associated with the risk of bloodstream infections (BSIs) and sepsis, which may be fatal. Therefore, continuous subcutaneous (SC) infusion is the preferred mode of administration.
- Do not abruptly lower the dose or withdraw dosing.
- Treprostinil injection may cause symptomatic hypotension.
- Titrate slowly in patients with hepatic insufficiency because such patients will likely be exposed to greater systemic concentrations relative to patients with normal hepatic function.
- Treprostinil injection inhibits platelet aggregation and increases the risk of bleeding.

ADVERSE REACTIONS

During clinical trials with SC infusion of treprostinil, infusion site pain and infusion site reaction (e.g., erythema, induration, or rash) were the most common adverse events and occurred in majority of those treated with treprostinil. Infusion site reactions were sometimes severe and led to discontinuation of treatment. Rash and hypotension (14% and 4%, respectively) were also commonly reported with SC infusion of treprostinil. Other common adverse events ($\geq 3\%$ more than placebo) included headache, diarrhea, jaw pain, edema, vasodilatation, and nausea, and these are generally considered to be related to the pharmacologic effects of treprostinil, whether administered subcutaneously or intravenously. The adverse reactions reported with treprostinil IV included bloodstream infections, arm swelling, paresthesia, hematoma, and pain.

DRUG INTERACTIONS

Treprostinil injection dosage adjustment may be necessary if inhibitors or inducers of CYP2C8 are added or withdrawn.

USE IN SPECIFIC POPULATIONS

- Safety and effectiveness of Treprostinil injection in pediatric patients have not been established.
- It is unknown if geriatric patients respond differently than younger patients. Caution should be used when selecting a dose for geriatric patients.
- There are no adequate and well-controlled studies with Treprostinil injection in pregnant women.
- It is not known whether Treprostinil injection is excreted in human milk.

Please see accompanying full Prescribing Information for additional safety information, also available by [clicking here](#).

Using this cover sheet, fax all pages of the enrollment form, along with the requested clinical documentation, to the Specialty Pharmacy below.

Date

TO

Accredo Health Group, Inc.

FAX 1-800-711-3526

Phone: 1-866-344-4874

FROM

(Name of agent of prescriber transmitting this fax/prescription)

Phone

Facility Name

Fax

RE

Patient Name

Date of Birth

DOCUMENTATION CHECKLIST

Indicate all current, signed and dated documents enclosed with this fax.

- | | |
|---|--|
| <input type="radio"/> Fully completed Treprostinil Enrollment Form, including: <ul style="list-style-type: none">- Patient/Insurance Information- Prescriber/Prescription Information- Medical Information/Patient Evaluation | <input type="radio"/> Echocardiogram |
| <input type="radio"/> Copy of front and back of Patient's Insurance card(s) | <input type="radio"/> 6-minute walk test results |
| <input type="radio"/> Right heart catheterization | <input type="radio"/> History and physical, including onset of symptoms, PAH clinical signs and symptoms and course of illness |
| | <input type="radio"/> Need for specific drug therapy |

Comments:

Number of Pages (including this cover sheet)